



Element concentrations in pelagic *Sargassum* along the Mexican Caribbean coast in 2018-2019

Rosa E. Rodríguez-Martínez^{1,*}, Priyadarsi D. Roy^{2,*}, Nuria Torrescano-Valle³, Nancy Cabanillas-Terán^{3,4}, Silvia Carrillo-Domínguez⁵, Ligia Collado-Vides⁶, Marta García-Sánchez^{1,7} and Brigitta I. van Tussenbroek¹

¹ Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Puerto Morelos, Quintana Roo, México

² Instituto de Geología, Universidad Nacional Autónoma de México, Ciudad de México, Ciudad de México, México

³ El Colegio de la Frontera Sur, Unidad Chetumal, Chetumal, Quintana Roo, México

⁴ CONACYT - El Colegio de la Frontera Sur, Chetumal, Quintana Roo, México

⁵ Departamento de Nutrición Animal, Instituto Nacional de Ciencias Médicas y Nutrición “Salvador Zubirán”, Ciudad de México, Ciudad de México, México

⁶ Department of Biological Sciences and Center for Coastal Oceans Research in the Institute of Environment, Florida International University, Miami, FL, United States of America

⁷ Instituto de Ingeniería, Universidad Nacional Autónoma de México, Ciudad de México, Ciudad de México, México

* These authors contributed equally to this work.

ABSTRACT

The massive influx of pelagic *Sargassum* spp. (sargasso) into the Mexican Caribbean Sea has caused major deterioration of the coastal environment and has affected the tourism industry as well as livelihoods since 2015. Species of *Sargassum* have high capacity to absorb metals; thus, leachates of sargasso may contribute to contamination by potentially toxic metals when they drain into the sea and into the groundwater when dumped in inadequate land deposits. Valorization of sargasso would contribute to sustainable management; therefore, knowledge on potentially toxic metal content is necessary to define possible uses of the algae. We present concentrations of 28 elements measured using a non-destructive X-ray fluorescence analyzer (XRF) in 63 samples of sargasso collected between August 2018 and June 2019 from eight localities along ~370 km long coastline of the Mexican Caribbean Sea. The sargasso tissues contained detectable concentrations of Al, As, Ca, Cl, Cu, Fe, K, Mg, Mn, Mo, P, Pb, Rb, S, Si, Sr, Th, U, V, and Zn. The element concentration in sargasso varied on spatial and temporal scales, which likely depended on the previous trajectory of the pelagic masses, and whether these had (or had not) passed through contaminated areas. Total arsenic concentration varied between 24–172 ppm DW, exceeding the maximum limit for seaweed intended as animal fodder (40 ppm DW) in 86% of the samples. For valorization, we recommend analyses of metal contents as a mandatory practice or avoiding uses for nutritional purposes. The high arsenic content is also of concern for environmental contamination of the sea and aquifer.

Submitted 25 November 2019

Accepted 30 January 2020

Published 26 February 2020

Corresponding author

Rosa E. Rodríguez-Martínez,
rosaer@cmar.unam.mx

Academic editor

Todd Anderson

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.8667

© Copyright

2020 Rodríguez-Martínez et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Aquatic and Marine Chemistry, Environmental Contamination and Remediation, Environmental Impacts

Keywords *Sargassum fluitans*, *S. natans*, Metal content, Arsenic, Caribbean Sea

INTRODUCTION

The west coast of Africa and some eastern Caribbean islands received unusual large quantities of pelagic *Sargassum* spp. (*S. fluitans* (Boergesen) Boergesen and *S. natans* (Linnaeus) Gallion; hereafter named sargasso) for the first time in 2011 (Gower, Young & King, 2013). In subsequent years, the range of massive sargasso influx extended over the Atlantic Ocean and whole Caribbean Sea. Wang et al. (2019) reported more than 20 million metric tons of sargasso in the open ocean in the peak month of June 2018, when the Great Atlantic Sargasso Belt extended for 8,850 km in total length. Beaching of sargasso has caused havoc to the Caribbean coastal ecosystems. Leachates and particulate organic matter from stranded decaying algal masses depleted the oxygen in near shore waters and reduced visibility of the water column, causing mortality of near-shore seagrasses and fauna (van Tussenbroek et al., 2017; Rodríguez-Martínez et al., 2019). Onshore and near shore masses of sargasso interfered with the seaward journeys of the juvenile turtles (Maurer, De Neef & Stapleton, 2015), affected sea turtle nestings (Maurer, Stapleton & Layman, 2018) and altered the trophic structure of the sea urchin *Diadema antillarum* in coastal marine systems (Cabanillas-Terán et al., 2019). Massive beachings also enhanced beach erosion (van Tussenbroek et al., 2017). Coastal ecosystem-based tourist industry is one of the major sources of income for the Caribbean countries (Langin, 2018) and the potential socio-economic impacts of ecosystem degradation due to sargasso influx have yet to be assessed.

The Mexican Caribbean coast began receiving massive amounts of sargasso during the late 2014 and it reached a peak in September 2015, when in the northern section of the coast between Cancun and Puerto Morelos an average of $\sim 2,360 \text{ m}^3$ of algae (mixed with sand, seagrasses and other algae) arrived per km of coastline (Rodríguez-Martínez, van Tussenbroek & Jordán-Dahlgren, 2016). During 2016, and 2017, the influxes decreased, increasing again in 2018, when in the peak month May $\sim 8,793 \text{ m}^3 \text{ km}^{-1}$ of algae (mixed with sand, seagrasses and other algae) were removed from the same shore section (Rodríguez-Martínez et al., 2019). In the tourist beaches, the algae removed from the beach and sea have been disposed in areas that are not properly prepared to avoid leakage of the leachates into the aquifer. In addition, the cleaning efforts have not covered the whole coastline and thousands of tons of sargasso have accumulated annually along the Mexican Caribbean coast.

Like other brown algae, species of *Sargassum* (including the pelagic ones) have high capacity to absorb metals and other elements (Kuyucak & Volesky, 1988; Davis, Volesky & Vieira, 2000). This high absorption capacity is attributed to the unique mixture of polysaccharides, mainly alginates, in their cell walls (Fourest & Volesky, 1997). At present, the *Sargassum* spp. are used for different commercial end products, such as fertilizers (Milledge & Harvey, 2016), textiles, paper and drugs (Oyesiku & Egunyomi, 2014), as well

as in the production of biogas (Wang et al., 2018). They have also been increasingly used as food for animals and humans, and therefore the high concentrations of contaminants, including heavy metals, may pose potential health risks (Reis & Duarte, 2018). Therefore, it is mandatory to evaluate elemental concentrations to ensure that acceptable levels are maintained in terms of health regulations (e.g., Fourest & Volesky, 1997). Previous studies on metal contents in sargasso, were either based on limited number of samples collected mostly from a single locality (e.g., Nigeria Oyesiku & Egunyomi, 2014; Dominican Republic, Fernández et al., 2017) or in a single season (e.g., Addico & De Graft-Johnson, 2016). Hence, it is unclear how much the metal contents can vary in the algal tissues across sites and seasons and between species.

In this study, we estimate concentrations of 28 different elements in sargasso tissues collected from the Mexican Caribbean coast, covering a linear north-south distance of 370 km. We hypothesize that the elemental contents are variable both in time and space. The determinations of metals and other elements from this study provide an essential baseline data for adequate management and potential uses of sargasso.

SURVEY METHODOLOGY

Study sites

We collected 63 samples of sargasso along the Mexican Caribbean coast, from Contoy Island, at the northern extreme, to Xcalak in the south (Fig. 1). This region receives an average precipitation of $\sim 1,061 \text{ mm y}^{-1}$ and the sea-surface temperature (SST) ranges from 25.1–29.9 °C (Rodríguez-Martínez et al., 2010). The Yucatan Current, a major branch of the Caribbean Current, transports the pelagic algal masses parallel to the Mexican Caribbean coastline. Easterly trade-winds dominate this region during the summer and mild cold fronts occur during the winter season. Trade-winds transport the superficial waters towards the shore, importing the pelagic masses of sargasso towards the coast.

The coastal environment consists of beaches, rocky shores, seagrass beds, coral reefs, mangroves, jungle and underground rivers (Hernández-Arana et al., 2015). All these ecosystems provide services to the tourism industry, a crucial component of the regional economy (Spalding et al., 2017). In the karstic Yucatan peninsula, the freshwater aquifer and seawater are constantly interacting; especially near the coast (Hernández-Terrones et al., 2011; Hernández-Terrones et al., 2015). This region has no other major industries besides tourism. At present, this region has the highest number of hotel rooms in Mexico and the number of rooms has increased from 3,206 in 1975 to 100,986 in 2017 (SEDETUR, 2019). Similarly, the resident population grew almost 15-folds, from less than 100,000 in 1970 to 1,501,785 in 2015 (INEGI, 2015). This rapid urban development has caused coastal pollution through influx of nutrients (Carruthers, van Tussenbroek & Dennison, 2005; Hernández-Terrones et al., 2011; Baker, Rodríguez-Martínez & Fogel, 2013; van Tussenbroek et al., 2017), sewage (Metcalf et al., 2011), and some metals (e.g., Lead, see Whelan III, van Tussenbroek & Santos, 2011) into the coastal ecosystems.




Figure 1 **Sampling sites.** Location of the sampling sites of sargasso along the Mexican Caribbean coast between August 2018 and June 2019. Map produced in QGIS 2.18 (<http://www.qgis.org>) using the following data sources: National Geospatial-Intelligence Agency (base map, World Vector Shoreline Plus, 2004. <http://shoreline.noaa.gov/data/datasheets/wvs.html>). The location of survey sites was obtained from the present study. Data sources are open access under the Creative Commons License (CC BY 4.0).

Full-size DOI: [10.7717/peerj.8667/fig-1](https://doi.org/10.7717/peerj.8667/fig-1)

Methodology

Field collection

Samples were collected between August 2018 and June 2019 from eight different sites along the Mexican Caribbean coast (from north to south): (1) Contoy Island, (2) Blue

Table 1 Samples information. Number of samples collected at eight sites along the Mexican Caribbean coast during 2018–2019. Habitat refers to distance from coast, shore (2–20 m from coast) or ocean (>5 km from coast).

Locality	Habitat	Collection		Species/Morphotype				Total
		Year	Month	Sflu III	Snat I	Snat VIII	Sarg sp	
1 - Contoy Island	Ocean	2019	March				4	4
2 - Blue waters	Ocean	2018	August		1			1
3 - Puerto Morelos	Shore	2018	August	1	1	1		3
			September	1	1	1		3
			October	2	1	2		5
			December	1	1			2
		2019	February	2	2	2		6
		March	1	3	2		6	
4 - Cozumel	Ocean	2019	April	1		1		2
			August			1		1
			May	1	1	1		3
5 - Mahahual	Shore	2019	May	4	3			7
6 - Chinchorro	Shore	2019	May		1			1
7 - Xahuayxol	Shore	2019	April	1	2			
			May	2	1			
			June	3	3			
8 - Xcalak	Shore	2019	May	3	3			6
Total				24	24	11	4	63

Notes.

Sarg sp: *Sargassum* spp., Sflu III: *Sargassum fluitans* III, Snat I: *S. natans* I, Snat VIII: *S. natans* VIII.

waters, (3) Puerto Morelos, (4) Cozumel, (5) Mahahual, (6) Chinchorro, (7) Xahuayxol and 8) Xcalak (Fig. 1, Table 1). Fresh sargasso (golden color) thalli floating near the shore (2–20 m) and in the ocean (>5 km from shore) were collected manually and separated in species and morphotypes (*S. fluitans* III, *S. natans* I and *S. natans* VIII) in the laboratory following Schell, Goodwin & Siuda (2015), except for the samples of Contoy Island (CI). The samples collected from CI were frozen before separating the specimens by species and morphotypes, thus, we classified them as *Sargassum* spp. All the samples were placed in an oven for at least 48 h at 60 °C until completely dry. Special caution was taken to avoid contact between the algal samples and any metal object. Samples were shipped to the Institute of Geology of the National Autonomous University of Mexico for the analysis of element concentrations. We did not remove epibionts from the thalli and analyzed the chemical composition of the algae including attached organisms, as the main interest of this study was to determine the potential contamination hazards and uses of sargasso as collected from the sea, without any specific separation treatment. All surveys were conducted under permit PPD/DGOPA-116/14 granted by SAGARPA (Agriculture, Natural Resources and Fisheries Secretariat) to B.I. van Tussenbroek.

Elemental analysis

Concentrations of 28 different elements were measured in dry samples using a Niton FXL 950 energy dispersive X-ray fluorescence (XRF) containing a 50 kV X-ray tube of Ag and

equipped with a geometrically optimized large area drift deflector following [Quiroz-Jiménez & Roy \(2017\)](#). [Table S1](#) shows the limit of detection of these elements. The dried samples were processed in the laboratory using a non-destructive sample preparation technique. Approximately 5–7 dry g of each sample was placed in a plastic capsule that has a 4 μm thick polypropylene X-ray film on one side and the other side of the capsule was packed with synthetic flexible gauze. The samples were measured in the mining Cu/Zn mode and three different filters using the internal calibration curves previously generated by comparing the results of Niton FXL with a conventional XRF (e.g., [Quiroz-Jiménez & Roy, 2017](#)). The results are expressed in parts per million dry weight (ppm DW) after carrying out the analysis in five repetitions in each sample. We used two different geological reference materials (Es-2, organic rich argillite and Es-4, dolostone) for estimation of precision ([Kiipli et al., 2000](#)). Except for Mg, all other elements have relative standard deviation (RSD) between <1 and 5%. Mg concentrations show RSD of 26% and it is the least precise among all the analyzed elements. Some advantages of the XRF analysis compared to other methodologies are that small samples are required (~5 g), the results have high precision, and it is non-destructive, permitting the same sample to be reused for other studies. Also, it is less expensive and faster compared to the use of an ICP-MS. The relatively high limit of detection of XRF for some elements is a disadvantage, and some potentially toxic elements may have been present in low concentrations, but were not measured (e.g., Ni and Co). This technique measures concentrations independent of the chemical state of an element.

Data analyses

The median of the five readings per element of each sample was calculated and used for further analysis. For each element, the readings below the limit of detection (<LOD; [Table S1](#)) were substituted with $LOD/\sqrt{2}$ for calculation of summary statistics ([Celo & Dabek-Zlotorzynska, 2010](#)). Distributions (spread of data and the median values) of the fourteen most commonly found elements (e.g., Al, As, Ca, Cl, K, Mg, Mn, P, Rb, S, Si, Sr, Th and U) in sargasso tissue for each sampling locality are illustrated by dot plots. Differences in the concentration of elements among species and morphotypes were tested using non-parametric ANOVAs based on the Kruskal–Wallis rank procedure. We constructed a heatmap using the data from fourteen elements from Puerto Morelos (location 3, see [Fig. 1](#)) to visualize temporal differences in concentration of metals in seven different sampling periods between August 2018 and April 2019. Element concentration values were Z -score-transformed across sampling times and their values above and below the mean were used to generate the heatmap. The Z -value is a dimensionless quantity which is defined by the following equation ([Larsen & Marx, 1986](#)):

$$Z = (X - \mu)/\sigma$$

Where X represents an individual raw score that is to be standardized, σ is the standard deviation of the population, and μ is the mean of the population.

All analyses were done in R ([R Core Team, 2019](#)) using packages: dplyr ([Wickham et al., 2019](#)), ggplot2 ([Wickham, 2009](#)), gplots ([Warnes et al., 2009](#)), pgirmess ([Giraudoux, 2013](#)), reshape ([Wickham, 2018](#)), tidyr ([Wickham & Henry, 2017](#)), and RColorBrewer

Table 2 Element concentrations median and range. Element concentrations (ppm DW) of pelagic *Sargassum* spp. tissue collected from eight localities along the Mexican Caribbean coast between 2018 and 2019. The number of samples with readings above LOD are expressed in % of the total sample size ($n = 63$).

Element	LOD	Samples with readings above LOD (%)	Minimum	Maximum	Median
Al	140	58.7	<LOD	500	206
As	4	100	24	172	80
Ca	394	100	23,723	136,146	70,040
Cl	266	100	747	53,101	22,350
Cu	6	7.9	<LOD	540	<LOD
Fe	3	7.9	<LOD	11	<LOD
K	333	100	1,990	46,002	19,666
Mg	2,915	92.1	<LOD	13,662	6,537
Mn	13	100	40	139	71
Mo	1	7.9	<LOD	7	<LOD
P	145	100	228	401	327
Pb	2	7.9	<LOD	3	<LOD
Rb	1	100	30	143	56
S	199	100	9,462	24,773	14,363
Si	342	100	447	2,922	1,767
Sr	6	100	1,605	2,564	1,890
Th	1	100	5	23	10
U	4	100	11	48	23
V	3	28.6	<LOD	13	<LOD
Zn	5	12.7	<LOD	17	<LOD

Notes.

LOD, Limit of detection.

(*Neuwirth, 2011*) A reproducible record of all statistical analyses is available on GitHub (<https://github.com/rerodriguezmtz/ElementsSar>). This includes all underlying data and R code for all analyses.

RESULTS

The most frequent elements in sargasso tissues, detected in 100% of the samples, were As, Ca, Cl, K, Mn, P, Rb, S, Si, Sr, Th, and U. They were followed in frequency by Mg (92.1% of samples) and Al (58.7% of samples) (Table 2). Other elements were found in fewer samples and they had median concentrations below the LOD: V (28.6% of samples), Zn (12.7% of samples), and Cu, Fe, Mo and Pb, present in 7.9% of samples (Table 2). Ba, Cd, Co, Cr, Ni, Ti, Y, and Zr remained below the LOD in all the samples (See Table S1 for LOD values). Some elements showed more than 5-fold difference between their minimal and maximal concentrations (ppm DW). For example, Cl showed 71.1-fold difference, K exhibited 23.1-fold difference, As had 7.2-fold difference, Si showed 6.5-fold difference and Ca exhibited 5.7-fold difference between their minimum and maximum values (Table 2).

Concentrations of P, S and Sr showed the least inter-site variability and the concentrations of Al, As, Cl and K showed the most inter-site variability (Fig. 2).

Among the potentially toxic elements, only As (median contents of 24–172 ppm DW) and Mn (median contents of 40–139 ppm DW) were present in all the samples (Table 2). Of all samples, 86% presented As concentrations above the maximum allowable concentration for seaweeds to be used as animal fooder under European regulations (40 ppm DW; EU, 2019), and 100% of the samples were above the maximum allowable concentration for agricultural soils in Mexico (22 ppm DW; NOM-147-SEMARNAT-SSA1-2004). Approximately 5% of our samples showed Cu concentrations above maximum tolerable level of dietary minerals for sheep (25 ppm DW) and cattle (100 ppm DW) (McDowell, 1992). Other potentially toxic elements (e.g., Mo, Pb and Zn) were detected in only 8–13% of the samples and they had median concentrations below the toxic limits for agricultural soils (see Table 2 and Table S2).

Concentrations of As, Ca, Cl, K, Mn, Rb and Si varied significantly among sargasso species/morphotypes (Fig. 2, Kruskal–Wallis test, $p < 0.05$; Table 3). As, Cl, K and Rb were significantly higher in *Sargassum natans* VIII compared to *S. natans* I. The concentrations of Ca and Si were significantly lower in *S. natans* VIII than in *S. fluitans* III and *S. natans* I. Similarly, the concentration of Mn was higher in *S. natans* I compared to *S. fluitans* III and *S. natans* VIII (Table 3). Contents of Al, Mg, P, S, Sr, Th and U did not vary significantly among species and morphotypes (KW, $p > 0.05$; Table 3). We did not compare the concentrations of Cu, Fe, Mo, Pb and Zn statistically among the species/morphotypes as their medians remained <LOD.

The concentrations of fourteen different elements (i.e., Al, As, Ca, Cl, K, Mg, Mn, P, Rb, S, Si, Sr, Th and U) in sargasso collected at Puerto Morelos in seven different sampling periods, from August 2018 to April 2019, showed considerable variability (Fig. 3). This inconsistent pattern indicates absence of any seasonal tendency in the elemental concentrations.

DISCUSSION

The sargasso tissues from the Mexican Caribbean had more As, Cu and Mn and less Cd, Cr, Pb and Zn compared to the chemical compositions of the algae biomass from Nigeria, Ghana and Dominican Republic (Table 4). Most striking was the high variability of element concentrations detected both in space (different sites along the coast) and time (different sampling months). This variability is likely partially due to the pelagic nature of the sargasso, as a result of increased uptake when exposed to areas rich in metals. It is unlikely that heavy metals were absorbed in near-shore waters of the Mexican Caribbean because this area lacks these elements in high concentrations, due to absence of major industrial, mining or heavy agricultural activities in the region. In addition, the absorption of metals by *Sargassum thunbergii* under experimental conditions was only clearly noticeable after ≥ 3 d exposure (Wu et al., 2010), whereas the residence time of sargasso in near-shore Mexican waters is usually in the order of hours when it is transported from the Yucatan Current towards the shore. Thus, the sargasso tissues likely acquired the heavy and trace elements before entering the Mexican coastal waters. Different contaminants are released




Figure 2 Spatial variability in element concentrations. Concentration of fourteen most frequent elements (ppm algal DW) in tissues of sargasso collected at eight sites along the Mexican Caribbean coast in 2018–2019. Note differences in scale of the Y-axis. Each dot corresponds to the median of the five XRF readings per sample. Color of the dot represents the sargasso species/morphotype. The horizontal black lines correspond to the median for each site. The dotted blue line corresponds to the limit of detection of the XRF equipment. A, Aluminum; B, Arsenic; C, Calcium; D, Chlorine; E, Potassium; F, Magnesium; G, Manganese; H, Phosphorus; I, Rubidium; J, Sulphur; K, Silicon; L, Strontium; M, Thorium; N, Uranium. Figure 1 and Table 1 have the site and sample details.

Full-size DOI: [10.7717/peerj.8667/fig-2](https://doi.org/10.7717/peerj.8667/fig-2)

Table 3 Elements concentrations in sargasso morphotypes. Median and range (in parenthesis) of elements (ppm DW) in three sargasso species/-morphotypes collected from eight localities along the Mexican Caribbean coast in 2018–2019. *P* values show summary of statistical analyses using Kruskal–Wallis H test (bold if significant) and the last column shows results of multiple comparison test.

Element	a) <i>S. fluitans</i> III (<i>n</i> = 24)	b) <i>S. natans</i> I (<i>n</i> = 24)	c) <i>S. natans</i> VIII (<i>n</i> = 11)	<i>P</i>	Multiple comparison test
Al	221 (<LOD-392)	198 (<LOD-500)	<LOD (<LOD-327)	0.7341	
As	59 (34–172)	55 (32–172)	123 (24–145)	0.0213	c > b
Ca	76,727 (46,599–115,260)	81,965 (37,260–136,146)	43,289 (23,723–75,849)	0.0013	(a = b) > c
Cl	21,487 (1,831–46,502)	10,122 (747–46,485)	32,086 (2,279–53,101)	0.0111	c > b
K	19,466 (3,620–44,280)	14,309 (1,990–39,642)	32,900 (4,902–46,002)	0.0121	c > b
Mg	6,376 (2,062–12,325)	6,385 (2,062–12,949)	6,883 (2,062–13,662)	0.5990	
Mn	70 (51–112)	89 (52–139)	56 (40–135)	0.0019	b > (a = c)
P	336 (229–401)	328 (262–394)	300 (228–350)	0.0590	
Rb	60 (32–102)	51 (30–143)	67 (48–120)	0.0071	c > b
S	14,341 (11,328–24,773)	12,776 (9,462–21,170)	16,231 (12,449–19,500)	0.1370	
Si	1,861 (927–2,877)	2,095 (696–2,564)	1,049 (447–2,135)	0.0009	(a = b) > c
Sr	1,934 (1,641–2,395)	1,876 (1,605–2,564)	1,793 (1,633–2,362)	0.4375	
Th	10 (5–17)	8 (6–23)	9 (8–20)	0.3021	
U	22 (11–48)	23 (12–47)	27 (16–45)	0.2321	

Notes.

LOD, limit of detection.

into the ocean, some as point sources and others more continuous, in different parts across the North Equatorial Recirculation Region of the Atlantic Ocean (NERR) and the Wider Caribbean Region (as a result of long-range transport). [Fernandez, Singh & Jaffé \(2007\)](#) recognized the discharge of sewage, mineral extracts, fertilizer and pesticide used in the agricultural sector as the principal pollution sources. The pelagic masses of sargasso might have been exposed to these contaminants depending on its trajectory in the ocean. The metal sequestration also involves complex mechanisms of ion exchange, chelation, adsorption, and ion entrapment in polysaccharide networks of the algae ([Volesky & Holan, 1995](#)). This ion entrapment, in turn, depends on the affinity of some divalent metals to alginates ([Haug, 1961](#)), and pH of the seawater also influences absorption of metals ([Davis, Volesky & Vieira, 2000](#)). Alginates are often characterized by the proportion of mannuronic (M) and guluronic (G) acids present in the polymer (M:G ratio), which may vary among




Figure 3 Temporal variability in element concentrations. Variability in concentration of fourteen different elements (ppm DW) in sargasso collected at Puerto Morelos between August 2018 and April 2019. Z-score transformations were applied to values of each element across all the sampling periods and their intensities above and below the mean are represented on the heatmap by red and yellow colors, respectively, as shown on the color key bar.

Full-size DOI: [10.7717/peerj.8667/fig-3](https://doi.org/10.7717/peerj.8667/fig-3)

and within species. For example, Mn concentration was higher in *S. natans* I, whereas Ca and Si concentrations were higher in *S. fluitans* III and *S. natans* I, and the concentrations of As, Cl, K and Rb were higher in *S. natans* VIII than in *S. natans* I. Variations in the metal concentrations among the sargasso species and morphological forms may be explained by different concentrations in their tissues, but also by differences in calcifying epifauna, such as bryozoans, tube polychaeta, and crustose coralline algae (Weis, 1968; Huffard *et al.*, 2014). Large differences in concentrations of Si (447–2,922 ppm DW) could be explained by different abundance of diatoms and silicoflagellates present in the samples (Takahashi & Blackwelder, 1992).

Sargasso samples from the Mexican Caribbean coast contained essential macro-elements for plants, like Ca (23,723–136,146 ppm DW), K (1,990–46,002 ppm DW), Mg (<2,915–13,662 ppm DW), P (228–401 ppm DW) and S (9,462–24,773 ppm DW), in addition to various micro-elements. Similar properties have been found in other *Sargassum* spp., making them adequate as complementary fertilizers as they enhance growth, seed germination and photosynthesis of crop plants on mineral-depleted soils (Sathya *et al.*, 2010; Kumari, Kaur & Bhatnagar, 2013; El-Din, 2015). Some micro-elements found in sargasso from Mexico, such as Cu, Mn, Mo and Zn, are micronutrients in low concentrations, but they are potentially toxic when present in high concentrations. In

Table 4 Element concentrations in different studies. Comparison of element concentration in sargasso from the Mexican Caribbean coast and other studies in different parts of the world.

Element	Site (Year) Nigeria ^a (2012)	Dominican Republic ^b (2015)	Ghana ^c (2015)	Mexican Caribbean ^d (2018–2019)
Al		303–4,188		<140–517
As		14–42	13–54	24–172
Ba		7–17		<36
Ca		96,901–133,400		23,723–136,146
Cd		0.1–0.3	78–119	<2
Cl			61–1353	747–53,101
Co		0.4–1		<11
Cr		2–56		<8
Cu		2–12	24–36	<6–540
Fe	8,700 ± 280	20–655	1,209–5,910	<3–11
K	28,000 ± 740	2,208–33,602		1,990–46,002
Mg	42,750 ± 3,500	10,211–18,241		<2915–13,662
Mn		16–32		40–139
Mo		0.6–3		<1–7
Ni		10–33		<10
P	96,500 ± 21,200	761–1,145		228–401
Pb		1–2	105–335	<2–3
Rb		0.3–10		30–143
Si		23,883–55,776		447–2,922
Sr		1,162–1,437		1,605–2,564
Th		0.04–0.4		5–23
Ti		37–92		<29
U		0.2–0.7		11–48
V		1–3		<3–13
Y	40 ± 0.0	0.1–0.8		<1
Zn	50 ± 0.0	13–21	16–100	<5–17
Zr		8–34		<2

Notes.^aOyesiku & Egunyomi, 2014 (mean and SD).^bFernández et al., 2017 (range).^cAddico & De Graft-Johnson, 2016 (range).^dThis study (range).

this study, we detected the presence of Cu (<8–540 ppm DW) and Mo (<1–7 ppm DW) in 7.9% of the samples, Zn (<2–17 ppm DW) in 12.7% of the samples and Mn (40–139 ppm DW) in all the samples. Cu concentrations exceeded safely limits recommended for agricultural soils by several countries in 5% of the samples (see Table S2). Similarly, about 8% of our samples contained Mo concentrations above the maximum level established for agricultural soils by Canada (i.e., 2 ppm DW), but these were below the limits established by Austria and Poland (i.e., 10 ppm DW). Mn content was above 100 ppm DW in 22% of the samples, considered toxic for some plant species, but acceptable for others that

can tolerate Mn up to 5,000 ppm DW (Howe, Malcolm & Dobson, 2004). Pb (<2–3 ppm DW) could be detected only in 7.9% of the samples, due to the limitation related to LOD of XRF analysis, and its concentration always remained below the toxic levels. Arsenic is of concern for the usages of sargasso as complementary fertilizer for crop plants. Limits of total As allowed for agricultural soils are between 15–50 ppm DW depending on the country (Table S2) (Belmonte-Serrato et al., 2010), thus, continuous application of sargasso (with total As between 24–172 ppm DW) may cause accumulation of As in the soils above allowable levels. High concentrations of As in soil may be toxic for the plants themselves, as it interferes with photosynthesis and other metabolic processes (Påhlsson, 1989; Ruiz Huerta & Armienta Hernández, 2012).

Sargasso could also be considered as animal fodder due to the presence of micro- and macro-elements, in addition to proteins, fibers and other components (Marín et al., 2009; Carrillo et al., 2012). However, approximately 86% of the samples had total As concentrations above the maximum level (40 ppm DW) allowable in Europe for animal feed materials derived from seaweed (EU, European Union). The toxicity of As depends on its chemical form, with inorganic As (trivalent state As III and pentavalent state As V) considered toxic (e.g., Yuan et al., 2007, Circuncisão et al., 2018), thus, even if total As concentrations are below 40 ppm DW, it is recommendable to carry out As speciation studies before using sargasso as animal fodder.

The (occasional) high contents of potentially toxic metals in sargasso is also a serious threat for the environment. The Mexican Caribbean coast has already received millions of tons of algae since late 2014. This accumulation over time, in addition to eutrophication and organic matter accumulation (Carruthers, van Tussenbroek & Dennison, 2005; Hernández-Terrones et al., 2011; Baker, Rodríguez-Martínez & Fogel, 2013; van Tussenbroek et al., 2017), is also a potential source of metal contamination for this region, even though levels of some potentially toxic elements like Cu, Mo, Zn, Mn and Pb were low. The sargasso removed from Mexican Caribbean beaches is presently deposited at abandoned limestone quarries, near the coast, without any treatment. The Yucatan Peninsula has a highly porous karst aquifer that is the only source of freshwater in the region. The pollutants from near surface deposits can easily infiltrate into the aquifer causing accumulation of As and other potentially toxic metals in the groundwater. Considering that water from the aquifer flows into the ocean through underground rivers, all these metals and excessive nutrients will eventually reach the marine environment (Carruthers, van Tussenbroek & Dennison, 2005; Metcalfe et al., 2011; Baker, Rodríguez-Martínez & Fogel, 2013). Prevention and mitigation measures are urgently needed to ensure that the massive influx of sargasso does not harm the coastal ecosystems and the tourism-based economy of countries located in the vicinity of the Great Atlantic *Sargassum* belt, including the Mexican Caribbean. The analyses of different specimens collected over longer periods and from different locations is required to obtain reliable information about metal contents in tissues.

CONCLUSION

In countries affected by the Great Atlantic *Sargassum* belt, the accumulation of decomposing sargasso on shores has harmed the coastal ecosystems, tourism-based economy and general

human well-being. The Mexican Caribbean coast has received millions of tons of sargasso since late 2014, and our study concludes that the massive influx might contribute with potentially toxic elements to the coastal ecosystems, including the aquifer. We observed relatively higher values of As, Cu and Mn and lower values of Cd, Cr and Pb compared to similar studies in countries affected by the *Sargassum* belt. Cu, Mo, Zn, Mn and Pb were present in lower contents but their accumulation over time might be a potential source of contamination in this region. Total arsenic in most samples exceeded the limit established for usage as animal fodder in Europe and for agricultural soil in several countries. Further studies on As speciation are required before using sargasso in food industries to determine if it complies with guidelines of international institutions and organizations (i.e., FAO, WHO). Chemical analysis should also be conducted using other methodologies such as an ICP-MS, with better limit of detection, before evaluating sargasso usages in food, pharmaceutical and agricultural industries. Governments and industries have the financial strengths, as well as the moral and legal responsibilities, to carry out regular analyses of specimens collected over long periods and from different locations required for obtaining reliable information about metal contents in the tissues of sargasso due to its unpredictable variability.

ACKNOWLEDGEMENTS

Special thanks to Elisa Vera Vázquez, for collecting sargasso samples from Puerto Morelos and Manta México Caribe A.C. for providing samples from Blue Waters and Isla Contoy. We also want to thank CEMIE-Oceano for the samples at Cozumel. Gabriela González López provided logistic help during fieldwork and in laboratory and Irma Gabriela Vargas-Martinez helped in XRF analysis. The authors claim no conflict of interest with this work.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The authors received no funding for this work.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Rosa E. Rodríguez-Martínez conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Priyadarsi D. Roy, Nuria Torrescano-Valle and Nancy Cabanillas-Terán conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

- Silvia Carrillo-Domínguez, Ligia Collado-Vides, Marta García-Sánchez and Brigitta I. van Tussenbroek conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

All surveys were conducted under permit PPD/DGOPA-116/14 granted by SAGARPA (Agriculture, Natural Resources and Fisheries Secretariat) to Brigitta I. van Tussenbroek.

Data Availability

The following information was supplied regarding data availability:

Data is available at <https://github.com/rerodriguezmtz/ElementsSar>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8667#supplemental-information>.

REFERENCES

- Addico GND, De Graft-Johnson KAA. 2016.** Preliminary investigation into the chemical composition of the invasive brown seaweed *Sargassum* along the West Coast of Ghana. *African Journal of Biotechnology* 15:2184–2191 DOI 10.5897/AJB2015.15177.
- Baker DM, Rodríguez-Martínez RE, Fogel ML. 2013.** Tourism's nitrogen footprint on a Mesoamerican coral reef. *Coral Reefs* 32:691–699 DOI 10.1007/s00338-013-1040-2.
- Belmonte-Serrato F, Romero-Díaz A, Alonso-Sarría F, Moreno-Brotóns J, Rojo-López S. 2010.** Afección de suelos agrícolas por metales pesados en áreas limítrofes a explotaciones mineras del sureste de España. *Papeles de Geografía* 51–52:45–54.
- Cabanillas-Terán N, Hernández-Arana HA, Ruiz-Zárate M-Á, Vega-Zepeda A, Sanchez-Gonzalez A. 2019.** *Sargassum* blooms in the Caribbean alter the trophic structure of the sea urchin *Diadema antillarum*. *PeerJ* 7:e7589 DOI 10.7717/peerj.7589.
- Carrillo S, Bahena A, Casas M, Carranco ME, Calvo CC, Ávila E, Pérez-Gil F. 2012.** The alga *Sargassum* spp. as alternative to reduce egg cholesterol content. *Cuban Journal of Agricultural Science* 46:181–186.
- Carruthers TJB, Van Tussenbroek BI, Dennison WC. 2005.** Influence of submarine springs and wastewater on nutrient dynamics of Caribbean seagrass meadows. *Estuarine, Coastal and Shelf Science* 64:191–199 DOI 10.1016/j.ecss.2005.01.015.
- Celo V, Dabek-Zlotorzynska E. 2010.** Concentration and source origin of trace metals in PM 2.5 collected at selected Canadian sites within the Canadian national air pollution surveillance program. In: *Urban airborne particulate matter*. Berlin, Heidelberg: Springer, 19–38 DOI 10.1007/978-3-642-12278-1_2.
- Circuncisão AR, Catarino MD, Cardoso SM, Silva AMS. 2018.** Minerals from macroalgae origin: health benefits and risks for consumers. *Marine Drugs* 16:1–30 DOI 10.3390/md16110400.

- Davis TA, Volesky B, Vieira RH SF. 2000.** *Sargassum* seaweed as biosorbent for heavy metals. *Water Research* **34**:4270–4278 DOI [10.1016/S0043-1354\(00\)00177-9](https://doi.org/10.1016/S0043-1354(00)00177-9).
- El-Din SMM. 2015.** Utilization of seaweed extracts as bio-fertilizers to stimulate the growth of wheat seedlings. *The Egyptian Journal of Experimental Biology* **11**:31–39.
- EU (European Union). 2019.** Commission Regulation (EU) 2019/1869 of 7 2019 amending and correcting Annex I to Directive 2002/32/EC of the European Parliament and of the Council as regards maximum levels for certain undesirable substances in animal feed. Available at <http://eur-lex.europa.eu/eli/reg/2019/1869/oj> (accessed on 29 December 2019).
- Fernández F, Boluda CJ, Olivera J, Guillermo LA. 2017.** Análisis elemental prospectivo de la biomasa algal acumulada en las costas de la República Dominicana durante 2015. *Centro Azúcar* **44**:11–22.
- Fernandez A, Singh A, Jaffé R. 2007.** A literature review on trace metals and organic compounds of anthropogenic origin in the Wider Caribbean Region. *Marine Pollution Bulletin* **54**:1681–1691 DOI [10.1016/j.marpolbul.2007.08.007](https://doi.org/10.1016/j.marpolbul.2007.08.007).
- Fourest E, Volesky B. 1997.** Alginate properties and heavy metal biosorption by marine algae. *Applied Biochemistry and Biotechnology* **67**:215–226 DOI [10.1007/BF02788799](https://doi.org/10.1007/BF02788799).
- Giraudoux P. 2013.** Pgirmess: data analysis in ecology. R package version 1.5.7. Available at <http://CRAN.R-project.org/package=pgirmess>.
- Gower J, Young E, King S. 2013.** Satellite images suggest a new *Sargassum* source region in 2011. *Remote Sensing Letters* **4**:764–773 DOI [10.1080/2150704X.2013.796433](https://doi.org/10.1080/2150704X.2013.796433).
- Haug A. 1961.** Affinity of some divalent metals to different types of alginates. *Acta Chemica Scandinavica* **15**:1794–1795 DOI [10.3891/acta.chem.scand.15-1794](https://doi.org/10.3891/acta.chem.scand.15-1794).
- Hernández-Arana HA, Vega-Zepeda A, Ruíz-Zárate MA, Falcón-Álvarez LI, López-Adame H, Herrera-Silveira J, Kaster J. 2015.** Transverse coastal corridor: from freshwater lakes to coral reefs ecosystems. In: Islebe GA, Calmé S, León Cortés J, Schmook B, eds. *Biodiversity and conservation of the Yucatán Peninsula*. Cham: Springer, 355–376.
- Hernández-Terrones LM, Null KA, Ortega-Camacho D, Paytan A. 2015.** Water quality assessment in the Mexican Caribbean: impacts on the coastal ecosystem. *Continental Shelf Research* **102**:62–72 DOI [10.1016/j.csr.2015.04.015](https://doi.org/10.1016/j.csr.2015.04.015).
- Hernández-Terrones L, Rebolledo-Vieyra M, Merino-Ibarra M, Soto M, Le-Cossec A, Monroy-Ríos E. 2011.** Groundwater pollution in a karstic region (NE Yucatan): baseline nutrient content and flux to coastal ecosystems. *Water, Air, & Soil Pollution* **218**:517–528 DOI [10.1007/s11270-010-0664-x](https://doi.org/10.1007/s11270-010-0664-x).
- Howe P, Malcolm H, Dobson S. 2004.** *Manganese and its components: environmental aspects*. Geneva: World Health Organization.
- Huffard CL, von Thun S, Sherman AD, Sealy K, Smith Jr KL. 2014.** Pelagic *Sargassum* community change over a 40-year period: temporal and spatial variability. *Marine Biology* **161**:2735–2751 DOI [10.1007/s00227-014-2539-y](https://doi.org/10.1007/s00227-014-2539-y).
- INEGI. 2015.** Encuesta Intercensal 2015. Tabulados Básicos. Quintana Roo, México. INEGI Available at <https://www.inegi.org.mx/programas/intercensal/2015/default.html#Tabulados>. (accessed on 11 November 2019).

- Kiipli T, Batchelor RA, Bernal JP, Cowing C, Hagel-Brunnstrom M, Ingham MN, Johnson D, Kivisilla J, Knaack CH, Kump P, Lozano R, Michiels D, Orlova K, Pirrus E, Rousseau RM, Ruzicka J, Sandström H, Willis JP. 2000. Seven sedimentary rock reference samples from Estonia. *Oil Shale* 17:215–224.
- Kumari R, Kaur I, Bhatnagar AK. 2013. Enhancing soil health and productivity of *Lycopersicon esculentum* Mill, using *Sargassum johnstonii* Setchell & Gardner as a soil conditioner and fertilizer. *Journal of Applied Phycology* 25:1225–1235 DOI 10.1007/s10811-012-9933-y.
- Kuyucak N, Volesky B. 1988. Biosorbents for recovery of metals from industrial solutions. *Biotechnology Letters* 10:137–142 DOI 10.1007/BF01024641.
- Langin K. 2018. Seaweed masses assault Caribbean islands. *Science* 360:1157–1158 DOI 10.1126/science.360.6394.1157.
- Larsen RJ, Marx ML. 1986. *An introduction to mathematical statistics and its applications*. New Jersey: Prentice-Hall Englewood Cliffs.
- Marín A, Casas-Valdez M, Carrillo S, Hernández H, Monroy A, Sanginés L, Pérez-Gil F. 2009. The marine algae *Sargassum* spp. (Sargassaceae) as feed for sheep in tropical and subtropical regions. *Revista de Biología Tropical* 57:1271–1281.
- Maurer AS, De Neef E, Stapleton S. 2015. *Sargassum* accumulation may spell trouble for nesting sea turtles. *Frontiers in Ecology and the Environment* 13:394–395 DOI 10.1890/1540-9295-13.7.394.
- Maurer AS, Stapleton SP, Layman CA. 2018. Impacts of the Caribbean *Sargassum* influx on sea turtle nesting. In: *Proceedings of the 71st Gulf and Caribbean Fisheries Institute*. 327–329.
- McDowell LR. 1992. *Minerals in animal and human nutrition*. San Diego: Academic Press Inc, 524.
- Metcalfe CD, Beddows PA, Gold BG, Metcalfe TL, Li H, Van Lavieren H. 2011. Contaminants in the coastal karst aquifer system along the Caribbean coast of the Yucatan Peninsula, Mexico. *Environmental Pollution* 159:991–997 DOI 10.1016/j.envpol.2010.11.031.
- Milledge JJ, Harvey PJ. 2016. Golden tides: problem or golden opportunity? The valorisation of *Sargassum* from beach inundations. *Journal of Marine Science and Engineering* 4:1–19 DOI 10.3390/jmse4030060.
- Neuwirth E. 2011. R package. Version 1.0-5. Available at <http://CRAN.R-project.org/package=RColorBrewer>.
- Oyesiku OO, Egunyomi A. 2014. Identification and chemical studies of pelagic masses of *Sargassum natans* (Linnaeus) Gaillon and *S. fluitans* (Borgessen) Borgesen (brown algae), found offshore in Ondo State, Nigeria. *African Journal of Biotechnology* 13:1188–1193 DOI 10.5897/AJB2013.12335.
- Påhlsson A-MB. 1989. Toxicity of heavy metals (Zn, Cu, Cd, Pb) to vascular plants. *Water, Air, & Soil Pollution* 47:287–319 DOI 10.1007/BF00279329.
- Quiroz-Jiménez JD, Roy PD. 2017. Evaluation of geochemical data by two different XRF spectrometers in sediments from the Santiaguillo Basin (state of Durango, Mexico). *Geofísica International* 57:305–315.

- R Core Team.** 2019. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <https://www.R-project.org/>.
- Reis VAT, Duarte AC.** 2018. Analytical methodologies for arsenic speciation in macroalgae: a critical review. *Trends in Analytical Chemistry* **102**:170–184 DOI [10.1016/j.trac.2018.02.003](https://doi.org/10.1016/j.trac.2018.02.003).
- Rodríguez-Martínez RE, Medina-Valmaseda AE, Blanchon P, Monroy-Velázquez LV, Almazán-Becerril A, Delgado-Pech B, Vásquez-Yeomans L, Francisco V, García-Rivas MC.** 2019. Faunal mortality associated with massive beaching and decomposition of pelagic Sargassum. *Marine Pollution Bulletin* **146**:201–205 DOI [10.1016/j.marpolbul.2019.06.015](https://doi.org/10.1016/j.marpolbul.2019.06.015).
- Rodríguez-Martínez RE, Ruíz-Rentería F, Van Tussenbroek B, Barba-Santos G, Escalante-Mancera E, Jordán-Garza G, Jordán-Dahlgren E.** 2010. Environmental state and tendencies of the Puerto Morelos CARICOMP site, Mexico. *Revista de Biología Tropical* **58**:23–43.
- Rodríguez-Martínez RE, Van Tussenbroek B, Jordán-Dahlgren E.** 2016. Afluencia masiva de sargazo pelágico a la costa del Caribe mexicano (2014–2015). In: Mendoza E, Quijano-Scheggia SI, Olivos-Ortiz A, Núñez Vázquez EJ, eds. *Florecimientos Algales Nocivos en México*. Ensenada: CICESE, 352–365.
- Ruiz Huerta EA, Armienta Hernández MA.** 2012. Acumulación de arsénico y metales pesados en maíz en suelos cercanos a jales o residuos mineros. *Revista Internacional de Contaminación Ambiental* **28**:103–117.
- Sathya B, Indu H, Seenivasan R, Geetha S.** 2010. Influence of seaweed liquid fertilizer on the growth and biochemical composition of legume crop, *Cajanus cajan* (L.) Mill sp. *Journal of Phytology* **2**:50–63.
- Schell JM, Goodwin DS, Siuda ANS.** 2015. Recent *Sargassum* inundation events in the Caribbean: shipboard observations reveal dominance of a previously rare form. *Oceanography* **28**:8–11.
- SEDETUR.** 2019. Indicadores Turísticos de Quintana Roo. México. Available at <https://qroo.gob.mx/sedetur/indicadores-turisticos> (accessed on 8 October 2019).
- Spalding M, Burke L, Wood SA, Ashpole J, Hutchison J, Zu Ermgassen P.** 2017. Mapping the global value and distribution of coral reef tourism. *Marine Policy* **82**:104–113 DOI [10.1016/j.marpol.2017.05.014](https://doi.org/10.1016/j.marpol.2017.05.014).
- Takahashi K, Blackwelder PL.** 1992. The spatial distribution of silicoflagellates in the region of the Gulf Stream warm-core ring 82B: application to water mass tracer studies. *Deep Sea Research Part A. Oceanographic Research Papers* **39**:S327–S346 DOI [10.1016/S0198-0149\(11\)80018-X](https://doi.org/10.1016/S0198-0149(11)80018-X).
- van Tussenbroek BI, Hernández-Arana HA, Rodríguez-Martínez RE, Espinoza-Avalos J, Canizales-Flores HM, González-Godoy CE, Barba-Santos MG, Vega-Zepeda A, Collado-Vides L.** 2017. Severe impacts of brown tides caused by *Sargassum* spp. on near-shore Caribbean seagrass communities. *Marine Pollution Bulletin* **122**:272–281 DOI [10.1016/j.marpolbul.2017.06.057](https://doi.org/10.1016/j.marpolbul.2017.06.057).
- Volesky B, Holan ZR.** 1995. Biosorption of heavy metals. *Biotechnology Progress* **11**:235–250 DOI [10.1021/bp00033a001](https://doi.org/10.1021/bp00033a001).

- Wang M, Hu C, Barnes BB, Mitchum G, Lapointe B, Montoya JP. 2019. The great Atlantic *Sargassum* belt. *Science* 365:83–87 DOI 10.1126/science.aaw7912.
- Wang S, Cao B, Abomohra AE-F, Hu Y, Wang Q, He Z, Xu S, Feng Y, Bernard UB, Jiang X. 2018. Comparative study of combustion properties of two seaweeds in a batch fluidized bed. *Combustion Science and Technology* 190:755–769 DOI 10.1080/00102202.2017.1407761.
- Warnes GR, Bolker B, Bonebakker L, Gentleman R. 2009. gplots: Various R programming tools for plotting data. R package version. Available at <https://cran.r-project.org/web/packages/gplots/index.html>.
- Weis JS. 1968. Fauna associated with pelagic *Sargassum* in the Gulf Stream. *The American Midland Naturalist Journal* 80:554–558 DOI 10.2307/2423550.
- Whelan III T, van Tussenbroek BI, Santos MGB. 2011. Changes in trace metals in *Thalassia testudinum* after hurricane impacts. *Marine Pollution Bulletin* 62:2797–2802 DOI 10.1016/j.marpolbul.2011.09.007.
- Wickham H. 2009. Elegant graphics for data analysis (ggplot2). Available at <http://had.co.nz/ggplot2/book>.
- Wickham H. 2018. Reshape. Version 0.8.8. Available at <https://CRAN.R-project.org/package=reshape>.
- Wickham H, François R, Henry L, Müller K. 2019. dplyr package. Version 0.8.3. Available at <https://cran.r-project.org/web/packages/dplyr/index.html>.
- Wickham H, Henry L. 2017. Tidy: easily tidy data with 'spread ()' and 'gather ()' functions. R package version 0.6.1. Available at <https://cran.r-project.org/web/packages/tidy/index.html>.
- Wu H, Zhan D, Liu H, Ding G, Liu W, Li M. 2010. Study on accumulation and degradation of heavy metals by the brown alga *Sargassum thunbergii*. *Marine Sciences* 34:69–74.
- Yuan CG, He B, Gao EL, Lü JX, Jiang GB. 2007. Evaluation of extraction methods for arsenic speciation in polluted soil and rotten ore by HPLC-HG-AFS analysis. *Microchimica Acta* 159:175–182 DOI 10.1007/s00604-006-0709-4.